



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

whence,  $\sin a = 2bc \sqrt{[s(s-1/a)(s-1/b)(s-1/c)]}$ ;

$$x = a \sin a = 2abc \sqrt{[s(s-1/a)(s-1/b)(s-1/c)]}$$

$$= \frac{1}{2abc} \sqrt{[(bc+ac+ab)(-bc+ac+ab)(bc-ac+ab)(bc+ac-ab)]}.$$

Solved similarly by W. S. Risley, H. C. Feemster, A. M. Harding, G. W. Hartwell, Elmer Schuyler, and A. H. Holmes.


### GEOMETRY.

403. Proposed by C. E. GITHENS, Wheeling, West Virginia.

In a triangular field the sides enclosing an obtuse angle are 35 rods and 48 rods in length. Two straight lines are drawn from this vertex, and are at right angles to these sides. If these lines intersect the base 16 rods apart, how long is the third side of the field?

Solution by J. SCHETFER, A. M., Hagerstown, Maryland.

Let there be two intersecting circles;  $AB$  the line of centers,  $C$  one point of intersection. Draw  $AC$ ,  $BC$ ,  $CD$ ,  $CE$ ; put  $AC=b$ ,  $BC=a$ ,  $DE=m$ , radius of circles  $O$  and  $O'$ , respectively,  $R$  and  $r$ .



In triangles  $ACE$  and  $BDC$ , respectively, we have  $2R-m : b = \cos(A+B) : \cos B$ ;  $2r-m : a = \cos(A+B) : \cos A$ . Eliminating  $\cos(A+B)$  and considering  $2R \cos A = b$ ,  $2r \cos B = a$ , we get  $a^2 R(2R-m) = b^2 r(2r-m)$  .. (1).

Since  $\sin A : \sin B = a : b$ , we have  $b^2(1-\cos^2 A) = a^2(1-\cos^2 B)$ ; therefore,  $b^2 r^2(4R^2 - b^2) = a^2 R^2(4r^2 - a^2)$ .

From (2) we get

$r^2 = \frac{a^4 R^2}{4R^2(a^2 - b^2) + b^4} \dots (3)$ . Substituting in (1) and making all necessary reductions, involving merely simple algebraic operations, we get

$$16(a^2 - b^2)(R^5 - 16m(a^2 - b^2)R^4 - 4(a^2 - b^2)(2b^2 - m^2)R^3 + 4a^2 b^2 m R^2 + b^4(a^2 - b^2 - 2m^2)R - b^6 m) = 0,$$

or, divided by  $16(a^2 - b^2)$ ,

$$R^5 - mR^4 - \frac{2b^2 - m^2}{4} R^3 + \frac{a^2 b^2 m}{4(a^2 - b^2)} R^2 + \frac{b^4(a^2 - b^2 - 2m^2)}{16(a^2 - b^2)} R - \frac{b^6 m}{16(a^2 - b^2)} = 0.$$

This equation of the fifth degree is to be solved for  $R$ , then  $r$  is found from (3), and  $AB = 2(R+r) - m$ . For  $a=35$ ,  $b=48$ ,  $m=16$ , the equation is

$$R^5 - 16R^4 - 1088R^3 - 10463.024R^2 + 488494.5467R + 6415370.2 = 0.$$